## LEMON FRESH 20

PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Wear protective eyewear (goggles, face shield, or safety glasses). Remove contaminated clothing and wash before reuse. Avoid contact with skin. Wash thoroughly with soap and water after handling.

#### FIRST AID

- · Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- · Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eve

#### If on skin or clothing:

- Take off contaminated clothing.
- · Rinse skin immediately with plenty of water for 15-20 minutes.

# f swallowed:

- Call poison control center or doctor immediately for treatment advice
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poiso control center or doctor.
- Do not give anything by mouth to an unconscious person

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going or treatment

ence of a moderate amount of organic soil (5% serum). Painted surfaces, Plastic (such as polystyrene or polypro-Disinfects, cleans, and deodorizes in one labor saving pylene), Sealed stone, Stainless steel, Vinyl and Plastic step. Effective against the following pathogens: (Bacteria:) Upholstery, Finished Woodwork. Pseudomonas aeruginosa (Pseudomonas), Staphylococ- DILUTION: 1:64 — 2 OUNCES PER GALLON OF WATER cus aureus (Staph), Streptococcus pyogenes (Strep), Enterobacter aerogenes. Enterobacter cloacae, Escherichia coli, Klebsiella pneumoniae, Mycoplasma gallisepticum, It is a violation of Federal Law to use this product in a man-Proteus mirabilis. Proteus vulgaris. Salmonella cholerae- ner inconsistent with its labeling. suis, Serratia marcescens, Shigella flexneri, Shigella son- GENERAL USE DIRECTIONS nei, Staphylococcus aureus phage 80 & 81, Staphylococ- APPLICATION: For disinfection, remove heavy soil decus epidermidis, (Viruses:) \*Adenovirus Type 3, \*Herpes posits from surface. Then thoroughly wet surface with a simplex Type 1 (Herpes), \*HIV-1 (AIDS virus), \*Influenza solution of 2 ounces of the concentrate per gallon of wa-A/Texas (Influenza), \*Rubella (German Measles), \*Vac- ter. Apply with a cloth, mop, sponge, or coarse sprayer,

cinia (Pox Virus), (Fungi:) Aspergillus flavus, Aspergillus fumigatus, Aspergillus glaucus var. tonophilus, Candida albicans, and Trichophyton mentagrophytes (Athlete's

USF IN: Bathrooms, Homes (households), Kitchens, Dental offices, Hospitals, Medical Offices (Clinics), Nursing homes. Day care centers. Nurseries. Bars. Cafeterias. Convenience Stores, Federally inspected meat and poultry plants, Food processing plants, Food storage areas, Institutional kitchens, Restaurants, (USDA) (Federally) inspected food processing facilities, Athletic facilities, Barber shops, Business and office buildings, Colleges, Correctional facilities, Dressing rooms, Exercise facilities, Factories, Hotels, Institutional facilities, Institutions, Locker rooms, Motels, Prisons, Public facilities, Public rest rooms, Schools, Shower and bath areas, Salons (Beauty), Camp grounds, Play ground equipment, Airplanes, Airports, Boats, Buses, Campers, Cars, Emergency vehicles, Mobile homes, Ships, Taxis, Trailers, Trains, Transportation terminals,

Use on washable hard, nonporous surfaces of: Exterior surfaces of Appliances, Basins (empty), Bathroom (lavatory) fixtures, Bathtubs, Cabinets, Cages, Chairs, Coils and drain pans of air conditioning and refrigeration equipment and heat pumps, Conductive flooring, Counters (countertops), Desks, Doorknobs, Floors, Garbage cans, Highchairs, Exterior surfaces of Microwave ovens, Outdoor furniture, Exterior surfaces of Refrigerators, Refrigerated storage and display equipment. Showers. Shower stalls Sinks (bathroom, kitchen), Stoves, (stovetops), Tables (picnic tables), Telephones, Toilets, Toilet bowls, Toilet bowl surfaces, Tubs, glazed Tiles, urinals, Walls. Other hard nonporous surfaces made of: Formica®, Glazed ce-A multi-purpose, germicidal detergent effective in the pres- ramic, Glazed enameled surfaces, Glazed porcelain, Metal,

# (488 ppm Active Quat) DIRECTIONS FOR USE



P.O. BOX 7453 • BEAUMONT, TX 77726-7453 PHONE: (800) 346-0132 FAX: (409) 866-4424 WWW CHEMAXMII COM

# FRESH 20

# **CLEANER • DEODORIZER • DISINFECTANT**

\*Virucidal • Disinfectant Pseudomonacidal • Staphylocidal • Salmonellacidal Bactericidal • Fungicidal • Mildewstatic

ACTIVE INGREDIENTS	
Didecyl dimethyl ammonium chloride	.1.875%
n-Alkyl (C14 50%, C12 40%, C16 10%) dimethyl benzyl	
ammonium chloride	1.248%
OTHER INGREDIENTS	96.877%
TOTAL	00.000%

## **PRODUCT No. C 206**

KEEP OUT OF REACH OF CHILDREN WARNING:

See Additional Precautionary Statements on Side Panel

EPA REG. NO. 47371-37-41442

EPA EST. NO. 44446-TX-1

Not for sale for or use by homeowners or residential users NET CONTENTS: 1 Quart

or by soaking. For sprayer applications, use a coarse MILDEWSTATIC INSTRUCTIONS: Will effectively spray device. Spray 6-8 inches from surface, rub with control the growth of mold and mildew plus the odors a brush, cloth or sponge. Do not breathe spray. Let caused by them when applied to hard, non-porous solution remain on surface for a minimum of 10 min- surfaces such as walls, floors and table tops. Apply utes. Rinse or wipe dry with a clean cloth or sponge solution [2 ounces (1/4 cup) per gallon of water] with or allow to air dry

as countertops, exterior surfaces of appliances, tables fresh solution for each use. Repeat application weekly and stovetops with potable water before reuse. Do not or when growth reappears.

use on utensils, plassware and dishes

or visibly soiled.

This product is not to be used as a terminal sterilant/ \*Kills HIV-1 (AIDS virus) on precleaned, environhigh level disinfectant on any surface or instrument mental surfaces/objects previously soiled with that (1) is introduced directly into the human body, ei- blood/body fluids in healthcare settings or other setther in contact with the bloodstream or normally sterile tings in which there is an expected likelihood of soiling areas of the body, or (2) contacts intact mucous mem- of inanimate surfaces/objects with blood/body fluids. branes but which does not ordinarily penetrate the and in which the surfaces/objects likely to be soiled blood barrier or otherwise enter normally sterile areas, with blood/body fluids can be associated with the poof the body. This product may be used to preclean or tential for transmission of Human Immunodeficiency decontaminate critical or semi-critical medical devic- Virus Type 1 (HIV-1) (associated with AIDS). es prior to sterilization or high level disinfection.

USDA: For use in federally inspected meat and poultry CONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF plants as a floor and wall cleaner for use in all depart- SURFACES/ OBJECTS SOILED WITH BLOOD/BODY ments. Food products and packaging materials must FLUIDS: Personal Protection: Clean-up should always be removed from the room or carefully protected, be done wearing protective gloves, gowns, masks and All surfaces must be thoroughly rinsed with potable eye protection.

may require brushing. To disinfect, first preclean to remove heavy soil, then remove or expel over the inner trap the residual bowl water. Pour in 3 ounces of the diluted solution. Swab the bowl completely using Disposal of Infectious Materials: Blood, body fluids, a scrub brush or mop, making sure to get under rim.

cleaning materials and clothing should be autoclaved and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal, state, and local and disposed of according to federal and dispos Leave for 10 minutes or overnight, then flush.

DISINFECTION OF BARBER / BEAUTY / SALON IN- HEAVY-DUTY CLEANING INSTRUCTIONS: Use 4 to 6 STRUMENTS AND TOOLS: Tools (such as combs, oz. of this product per gallon of water to clean heavily brushes, razors, scissors, and other shop tools) can solled areas. solution for a contact time of 10 minutes. Prepare a Store in original container in areas inaccessible to chilcomes diluted or visibly soiled.

a cloth, mop, sponge or coarse sprayer, making sure Rinse all surfaces that come in contact with food such to wet all surfaces completely. Let air dry. Prepare a

FOR FUNGICIDAL ACTIVITY: This product is an effec-Rinsing of floors is not necessary unless they are tive fungicide against Trichophy-ton mentagrophytes to be waxed or polished. Prepare a fresh solution (the athlete's foot fungus) in areas such as locker daily or more often if the solution becomes diluted rooms, dressing rooms, shower and bath areas and exercise facilities. Follow disinfection directions.

SPECIAL INSTRUCTIONS FOR CLEANING AND DE-

Cleaning procedures: Blood and other body fluids TOILET BOWLS: To clean, apply diluted solution must be thoroughly cleaned from surfaces and objects around the bowl and up under the rim. Stubborn stains

regulations for infectious waste disposal.

fresh solution daily or more often if the solution beof side walls. Do not reuse empty container, Discard in trash or recycle.